

ON A TWO-DIMENSIONAL SYMPLECTIC SPACE-TIME

Joachim Nzotungicimpaye*
Kigali Institute of Education
Department of Mathematics
P.O.Box 5039
e-mail kimpaye @kie.ac.rw

February 7, 2008

Abstract

We realize the hamiltonian action of the one spatial Poincare group G , analogous of the projective unitary representation, on a coadjoint orbit of G which we interpret as a symplectic space-time where the time is associated to a physical quantity of cohomological origin (the force f). This time is the time delay necessary for the system under the force f to have momentum p .

1 Introduction

One can verify that the one dimensional Poincare group [1] (Galilei group [1]) G is a subgroup of the real inhomogeneous symplectic group $ISP(2, R)$. It can then be seen as a group of canonical transformations for a certain two dimensional symplectic manifold. In this paper, using the methods of symplectic realizations of Lie groups [5], we realize such a symplectic manifold as a space-time which is a coadjoint orbit of the Poincare group G on the dual $\mathcal{H}^*$ of the extension of its Lie algebra. By the contraction process [2], we also obtain the realization of the one spatial Galilei group on the same space-time. The paper is organized as follows. In section 2 we study the central extension H of G and the coadjoint action of G on $\mathcal{H}^*$. In the section 3 we realize the one spatial space-time as an coadjoint orbit of G and obtain the Poincare and the Galilei groups as canonical transformations of that space-time.

*Previously professor at the University of Burundi

2 Central extension of G and its coadjoint action on $\mathcal{H}^*$

First of all, let us recall that the general element of the one spatial Poincare group G is parametrized as $g = (v, \tau, x)$ where v is a Lorentz boost, τ is a time translation while x is a space translation. The multiplication law is [1]

$$(v, \tau, x)(v', \tau', x') = \left(\frac{v + v'}{1 + \frac{vv'}{c^2}}, \gamma\tau' + \gamma\frac{vx'}{c^2} + \tau, \gamma v\tau' + \gamma x' + x \right) \quad (1)$$

where

$$\gamma^{-1} = \sqrt{1 - \frac{v^2}{c^2}} \quad (2)$$

c being the velocity of light in a empty space. If K, P and E are respectively the left invariant vector fields generating the boosts, time translations and space translations, then the Poincare Lie algebra $\mathcal{G}$ has the structure

$$[K, E] = P, [K, P] = \frac{1}{c^2}E \quad (3)$$

One can verify [4] that the central extension $\mathcal{H}$ of this Lie algebra is generated by K, P, E and F such that the nontrivial Lie brackets are

$$[K, E] = P, [K, P] = \frac{1}{c^2}E, [P, E] = F \quad (4)$$

Now writing the general element h of the corresponding connected Lie group H as $h = \exp(\zeta F)\exp(xP + tE)\exp(vK)$, and using the Baker-Campbell-Hausdorff formulae, we find that the multiplication law for H is

$$\begin{aligned} & (v, \tau, x, \zeta)(v', \tau', x', \zeta') \\ &= \left(\frac{v + v'}{1 + \frac{vv'}{c^2}}, \gamma\tau' + \gamma\frac{vx'}{c^2} + \tau, \gamma v\tau' + \gamma x' + x, \zeta + \zeta' + \frac{1}{2}\gamma(x - v\tau)\tau' - \frac{1}{2}\gamma(\tau - \frac{vx}{c^2})x' \right) \end{aligned} \quad (5)$$

We then verify that the coadjoint action of G on $\mathcal{H}^*$ is

$$\begin{aligned} & Ad_{(v, \tau, x)}^*(k, e, p, f) \\ &= \left(k + \gamma(x - v\tau)\frac{e}{c^2} + \gamma(\tau - \frac{vx}{c^2})p + \frac{1}{2}(\tau^2 - \frac{x^2}{c^2})f, \gamma e - \gamma pv - fx, -\frac{\gamma v}{c^2}e + \gamma p + f\tau, f \right) \end{aligned} \quad (6)$$

3 Symplectic structure on space-time

Using (4), one finds that the Kirillov form [3], [5] is

$$\begin{pmatrix} 0 & -p & -\frac{e}{c^2} \\ p & 0 & f \\ \frac{e}{c^2} & -f & 0 \end{pmatrix}$$

and that the coadjoint orbit is characterized by the invariants f and

$$\mathcal{K} = k + f \frac{q^2}{2c^2} - f \frac{t^2}{2} \quad (7)$$

where $(t = \frac{p}{f}, q = -\frac{e}{f})$. We denote it by $\mathcal{O}_{(f,\mathcal{K})}$. The symplectic form is, in Darboux's coordinates,

$$\sigma = dp \wedge dq \quad (8)$$

By use of the result (6), the symplectic realization of the Poincare group is then given by

$$L_{(v,\tau,x)}(p, q) = (\gamma p + \gamma \frac{fv}{c^2} q + f\tau, \gamma \frac{v}{f} p + \gamma q + x) \quad (9)$$

It depends explicitly on invariant f . This explicit dependence can be removed if we use, instead of the Darboux's coordinates (p, q) , the space-time coordinates (q, t) . In that case, (9) becomes

$$L_{(v,\tau,x)}(t, q) = (\gamma t + \gamma \frac{v}{c^2} q + \tau, \gamma vt + \gamma q + x) \quad (10)$$

which are the usual Poincare transformations of space-time. The dependence on f is transferred to the symplectic form which become $\sigma = f dt \wedge dq$. One can also verify that

$$ds^2 = -dq^2 + c^2 dt^2 \quad (11)$$

is invariant. The orbit $\mathcal{O}_{(f,\mathcal{K})}$ is then endowed with two geometric structures: the symplectic one given by (7) and the pseudo-euclidean one given by (10). If one neglects $\frac{v}{c}$, the action (10) becomes the usual Galilei transformation of space-time while (9) becomes the symplectic realization of the Galilei group on $\mathcal{O}_{(f,\mathcal{K})}$ [3] which as we have seen represents a massless particle under the constant force f . The orbit $\mathcal{O}_{(f,\mathcal{K})}$ can then be interpreted as a Poincare (Galilei) elementary system, characterized by a position q , a linear momentum p under an invariant force f . The time t represents the time delay necessary to the system under the force f to have momentum p .

References

- [1] Levy-Leblond, J.M., *Ann. Ins. Henri Poincare*, 3, 1—, 1965
- [2] Inonu, E; Wigner, E.P., *Natl. Sci. USA*, 39, 510—, 1953
- [3] Nzotungicimpaye, J., *Letters in Mathematical Physics*, 15, 101—, 1988
- [4] Hamermesh, M., *Group theory and its applications to physical problems*, Addison Wesley, New York, 1962
- [5] Aldaya, V. and de Azcarraga, J.A., *Revista Nuovo Cimento*, 310, 1—, 1980
 Giachetti, *Revista Nuovo Cimento*, 412, 1—, 1981
 Matolsi, T., *Ann. Univ. Bud. sect. Math.* 20, 71—, 1977
 Souriau, J.M., *Structures des systmes dynamiques*, Dunod, Paris, 1970